



Publisher Correction: The clinical characteristics of patients with congenital nephrotic syndrome secondary to *NPHS1* mutation: Is nephrectomy still a therapeutic option for selected cases?

Yüksel Uğurlu¹ · Bora Gülhan² · İsmail Dursun³ · Hülya Nalçacıoğlu⁴ · Gülşah Kaya Aksoy⁵ · Nur Canpolat⁶ · Aysun Bayazıt⁷ · Zeynep Birsin Özçakar⁸ · Selcuk Yüksel⁹ · Gönül Parmaksız¹⁰ · Gülşah Özdemir² · Eda Didem Kurt-Şükür² · Ali Düzova² · Mutlu Hayran¹¹ · Fatih Ozaltın^{2,12,13,14}

Published online: 9 May 2025
© The Author(s) 2025

Publisher Correction: Pediatric Nephrology

<https://doi.org/10.1007/s00467-025-06774-6>

The legend for Fig. 1 was previously omitted and has now been included as follows:

"I: Nephrectomy; ✦: eGFR <60 ml/min/1.73 m²; ∇: Kidney transplantation; orange: albumin <2.5 g/dL; blue: albumin >2.5 g/dL; x: Exitus; three squares represent one year; Yes*: Sequential bilateral nephrectomy; Yes**: Concurrent bilateral nephrectomy."

The original article has been corrected, and the publisher apologizes for the inconvenience.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00467-025-06774-6>.

✉ Fatih Ozaltın
fozaltin@hacettepe.edu.tr

¹ Department of Pediatrics, Faculty of Medicine, Hacettepe University, Ankara, Türkiye

² Department of Pediatric Nephrology, Faculty of Medicine, Hacettepe University, Sıhhiye, 06100 Ankara, Türkiye

³ Department of Pediatric Nephrology, Faculty of Medicine, Erciyes University, Kayseri, Türkiye

⁴ Department of Pediatric Nephrology, Faculty of Medicine, Ondokuz Mayıs University, Samsun, Türkiye

⁵ Department of Pediatric Nephrology, Faculty of Medicine, Akdeniz University, Antalya, Türkiye

⁶ Department of Pediatric Nephrology, Istanbul University-Cerrahpasa, Cerrahpasa Faculty of Medicine, Istanbul, Türkiye

⁷ Department of Pediatric Nephrology, Faculty of Medicine, Çukurova University, Adana, Türkiye

⁸ Department of Pediatric Nephrology, Faculty of Medicine, Ankara University, Ankara, Türkiye

⁹ Department of Pediatric Nephrology, Faculty of Medicine, Çanakkale Onsekiz Mart University, Çanakkale, Türkiye

¹⁰ Department of Pediatric Nephrology, Başkent University Adana Dr. Turgut Noyan Training and Research Center, Adana, Türkiye

¹¹ Department of Preventive Oncology, Hacettepe University Cancer Institute, Ankara, Türkiye

¹² Nephrogenetics Laboratory, Department of Pediatric Nephrology, Faculty of Medicine, Hacettepe University, Ankara, Türkiye

¹³ Center for Genomics and Rare Diseases, Hacettepe University, Ankara, Türkiye

¹⁴ Department of Bioinformatics, Hacettepe University Institute of Health Sciences, Ankara, Türkiye