

# ANNALS OF ANIMAL SCIENCE

ISSN: 2300-8733, <https://sciendo.com/journal/AOAS>

ACCEPTED AUTHOR VERSION OF THE MANUSCRIPT:

## Impacts of crowding stress on aquatic animals and its mitigation through feed additives supplementation – a review

DOI: 10.2478/aoas-2025-0038

Lee Seong Wei<sup>1♦</sup>, Martina Irwan Khoo<sup>2♦</sup>, Ramasamy Harikrishnan<sup>3</sup>, Ümit Acar<sup>4,5</sup>, Md Eilious Hosain<sup>6</sup>, Mohamad Nor Azra<sup>7,8</sup>, Ahmad Syazni Kamarudin<sup>9</sup>, Lau Kia Kian<sup>10</sup>, Hadiana<sup>11♦</sup>, Wendy Wee<sup>12♦</sup>

<sup>1</sup>Department of Agricultural Sciences, Faculty of Agro-Based Industry, Universiti Malaysia Kelantan, Jeli Campus, 17600 Jeli, Kelantan, Malaysia

<sup>2</sup>Department of Chemical Pathology, School of Medical Sciences, Universiti Sains Malaysia, Health Campus, Kubang Kerian 16150, Kelantan, Malaysia

<sup>3</sup>Department of Zoology, Pachaiyappa's College for Men (affiliation to University of Madras), Kanchipuram 631 501, Tamil Nadu, India

<sup>4</sup>Çanakkale Onsekiz Mart University, Bayramiç Vocational School, Department of Forestry, Çanakkale, Türkiye

<sup>5</sup>Çanakkale Onsekiz Mart University, Faculty of Marine Sciences and Technology, Department of Fisheries Industry Engineering, Çanakkale, Türkiye

<sup>6</sup>Fisheries Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka, 1349, Bangladesh

<sup>7</sup>Institute of Climate Adaptation and Marine Biotechnology (ICAMB), Universiti Malaysia Terengganu (UMT), Kuala Nerus 21030, Terengganu, Malaysia

<sup>8</sup>Research Center for Marine and Land Bioindustry, Earth Sciences and Maritime Organization, National Research and Innovation Agency (BRIN), Pemenang 83352, Indonesia

<sup>9</sup>School of Animal Science, Aquatic Science and Environment, Faculty of Bioresources and Food Industry, Universiti Sultan Zainal Abidin, Besut Campus, 22200 Besut, Terengganu, Malaysia

<sup>10</sup>Faculty of Pharmaceutical Sciences, UCSI University Kuala Lumpur Campus, No. 1, Jalan UCSI, UCSI Height, Cheras, 56000, Kuala Lumpur, Malaysia

<sup>11</sup>PSDKU Aquaculture, Faculty of Fisheries and Marine Science, Brawijaya University. Jl. Pringgodani, Kel. Mrican, Kec. Mojoroto. Kediri City, East Java, Indonesia, 64111

<sup>12</sup>Center for Fundamental and Continuing Education, Universiti Malaysia Terengganu, 21030, Kuala Nerus, Terengganu, Malaysia

♦Corresponding author: leeseong@umk.edu.my; msirwankhoo@gmail.com; hadiana23@ub.ac.id; wendy@umt.edu.my

Received date: 28 October 2024

Accepted date: 11 March 2025

**To cite this article:** (2025). Wei L.S., Khoo M.I., Harikrishnan R., Acar Ü., Hosain M.E., Azra M.N., Kamarudin A.S., Kian L.K., Hadiana, Wee W. (2025). Impacts of crowding stress on aquatic animals and its mitigation through feed additives supplementation – a review, *Annals of Animal Science*, DOI: 10.2478/aoas-2025-0038

**This is unedited PDF of peer-reviewed and accepted manuscript. Copyediting, typesetting, and review of the manuscript may affect the content, so this provisional version can differ from the final version.**

## Impacts of crowding stress on aquatic animals and its mitigation through feed additives supplementation – a review

Lee Seong Wei<sup>1♦</sup>, Martina Irwan Khoo<sup>2♦</sup>, Ramasamy Harikrishnan<sup>3</sup>, Ümit Acar<sup>4,5</sup>, Md Eilious Hosain<sup>6</sup>, Mohamad Nor Azra<sup>7,8</sup>, Ahmad Syazni Kamarudin<sup>9</sup>, Lau Kia Kian<sup>10</sup>, Hadiana<sup>11♦</sup>, Wendy Wee<sup>12♦</sup>

<sup>1</sup>Department of Agricultural Sciences, Faculty of Agro-Based Industry, Universiti Malaysia Kelantan, Jeli Campus, 17600 Jeli, Kelantan, Malaysia

<sup>2</sup>Department of Chemical Pathology, School of Medical Sciences, Universiti Sains Malaysia, Health Campus, Kubang Kerian 16150, Kelantan, Malaysia

<sup>3</sup>Department of Zoology, Pachaiyappa's College for Men (affiliation to University of Madras), Kanchipuram 631 501, Tamil Nadu, India

<sup>4</sup>Çanakkale Onsekiz Mart University, Bayramiç Vocational School, Department of Forestry, Çanakkale, Türkiye

<sup>5</sup>Çanakkale Onsekiz Mart University, Faculty of Marine Sciences and Technology, Department of Fisheries Industry Engineering, Çanakkale, Türkiye

<sup>6</sup>Fisheries Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka, 1349, Bangladesh

<sup>7</sup>Institute of Climate Adaptation and Marine Biotechnology (ICAMB), Universiti Malaysia Terengganu (UMT), Kuala Nerus 21030, Terengganu, Malaysia

<sup>8</sup>Research Center for Marine and Land Bioindustry, Earth Sciences and Maritime Organization, National Research and Innovation Agency (BRIN), Pemenang 83352, Indonesia

<sup>9</sup>School of Animal Science, Aquatic Science and Environment, Faculty of Bioresources and Food Industry, Universiti Sultan Zainal Abidin, Besut Campus, 22200 Besut, Terengganu, Malaysia

<sup>10</sup>Faculty of Pharmaceutical Sciences, UCSI University Kuala Lumpur Campus, No. 1, Jalan UCSI, UCSI Height, Cheras, 56000, Kuala Lumpur, Malaysia

<sup>11</sup>PSDKU Aquaculture, Faculty of Fisheries and Marine Science, Brawijaya University. Jl. Pringgodani, Kel. Mrican, Kec. Mojojoto. Kediri City, East Java, Indonesia, 64111

<sup>12</sup>Center for Fundamental and Continuing Education, Universiti Malaysia Terengganu, 21030, Kuala Nerus, Terengganu, Malaysia

♦Corresponding authors: leeseong@umk.edu.my; msirwankhoo@gmail.com; hadiana23@ub.ac.id; wendy@umt.edu.my

DOI: 10.2478/aoas-2025-0038

### Abstract

Continuous and sustainable aquaculture development is essential to provide affordable protein sources for the global population. Nevertheless, the intensification of fish farming at high stocking density led to chronic crowding stress that could impair the growth performance, immune system, and antioxidant capacity of an aquaculture species. Consequently, the prevalence of disease outbreaks increases, resulting in low aquaculture production. Various

countries remain highly dependent on antibiotics and chemotherapies to overcome this problem. Recently, researchers have proposed and developed alternative solutions using environmentally-friendly feed additives, such as probiotics, phytobiotics, and synthetic immune stimulants, to mitigate crowding stress. The feed additives reportedly alleviate crowding stress, promote growth performance, enhance gut microbiota, immune system, and antioxidant capacity, and stimulate disease resistance, contributing to the overall improvement of the health status of aquaculture species and boosting aquaculture production. The application of feed additives in aquafeed has been investigated in various aquaculture species to explore their beneficial effects at optimum doses. The study outcomes illustrated that feed additives in aquafeed are promising in mitigating crowding stress among aquaculture species. Herein, this review paper discussed and summarised the impacts of crowding stress on the growth, immune system, and antioxidant capacity of aquatic animals. In addition, this review highlighted the potential of feed additives such as probiotics, phytobiotics, and synthetic immune stimulants in mitigating crowding stress in aquatic animals was highlighted in this review.

**Key words:** crowding stress, probiotic, phytobiotic, synthetic immune stimulant, growth performance, immune system, antioxidant

Water quality, nutritious feed, and genetics are crucial in ensuring the success of aquaculture. On top of that, stocking density could impact the welfare and productivity of an aquaculture species. A non-optimal stocking density will result in poor water quality due to ammonia and nitrite accumulation and dissolved oxygen depletion in an aquaculture system (Amano et al., 2022). Moreover, the aquaculture species will exhibit abnormal behaviour and appetite loss, thus reducing farm productivity (Amano et al., 2022). As aquaculture is gearing toward intensification, the risk of crowding stress has become a setback for aquaculture industry development (Lin et al., 2018; Liu et al., 2014). Previous studies reported that crowding stress significantly compromised the immune system of aquaculture species, exposing them to a higher risk of disease infection (H. Adineh et al., 2021; Lin et al., 2018; Paray et al., 2020).

Cortisol is a hormone for stress regulation in aquatic animals (Gabr et al., 2023). Fish under stress exhibit high blood cortisol and glucose levels, which impairs their growth, physiology, and immune system (Espelid et al., 1996; Esteban et al., 2004; Sadoul & Vijayan, 2016; Schreck, 2010; Schreck & Tort, 2016), thus, impacting the fish welfare. Fish welfare is the primary for a successful aquaculture activity; hence, fish farmers are required to practice good stress management for their aquaculture species. Nonetheless, farmers face challenges in preventing crowding stress among the aquatic species as the aquaculture industry intensifies. Therefore, several strategies were utilised to maximise production while maintaining good welfare for the aquaculture species, although farmers are still struggling to find the right balance between the two components. Besides improving the fish farming facilities, incorporating feed additives is a practical and cost-effective strategy to maintain fish welfare in intensive aquaculture farming (Herrera et al., 2019).

Feed additives application is currently trending among aquaculturists. Studies have demonstrated the efficacy of feed additives in aquatic animals associated with handling (Costas

et al., 2012), salinity (Do Huu et al., 2016), anoxia (Pahor-Filho et al., 2017), ammonia (Rama & Manjabhat, 2014), hypoxia (Henrique et al., 1998), temperature (Chen et al., 2004), pH (Wang et al., 2014), starvation (Yengkokpam et al., 2016) and crowding stress (Castro-Osses et al., 2017; Chen et al., 2016; Ganga et al., 2011). The benefits of feed additives in aquaculture include improvements in fish growth performance (Güroy et al., 2012; Hamid et al., 2022), nutritional profile (Anis Mohamad Sukri et al., 2022), reproduction (Dabrowski & Ciereszko, 2001), immune system (Sukri et al., 2023), disease resistance (Abdul Kari et al., 2022; Goh et al., 2023), and mitigating stressor impacts (Kari, Wee, Hamid, et al., 2022; Wei et al., 2022). A recent study revealed the potential of phytobiotics and probiotics as feed additives to mitigate crowding stress in aquaculture.

Phytobiotics contain bioactive compounds, such as tannins, flavonoids, carotenoids, phenolic compounds, and terpenes, that can reduce stress and modulate the immune system (Kari et al., 2024; Seong Wei et al., 2024; Tungmunnithum et al., 2018; Wei et al., 2022). Furthermore, phytobiotics could promote the growth of gut microbiota, enhance feed utilisation efficiency, and stimulate disease resistance in aquatic animals (Abdul Kari et al., 2022; Goh et al., 2023; Kari, Wee, Abdul Hamid et al., 2022; Kari, Wee, Hamid, et al., 2022; Wei et al., 2022). Meanwhile, probiotics reportedly enhanced the gut microbiota balance (Kiriara et al., 2018) and intestinal microflora (Gatesoupe, 1999), eliminate gut pathogens, and stimulate the innate immune system (Gómez & Balcázar, 2008; Wee et al., 2024). Examples of probiotics in aquaculture are *Bacillus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Leuconostoc*, *Enterococcus*, *Aeromonas*, *Shewanella*, and *Carnobacterium* (Takanashi et al., 2014). Synthetic immune stimulants (SIS) were also used as a feed additive in mitigating stress among aquatic animals, including gallic acid in *Cyprinus carpio* (Ghafarifarsani et al., 2023) and vitamin E and C in gilthead seabream, *Sparus aurata* (Montero et al., 1999).

Given the accumulating evidence of crowding stress impact on aquatic species, the present review discussed and summarised the detrimental effects of this condition on animal growth and their immune system. Additionally, this review highlighted the potential of feed additives in mitigating crowding stress in aquaculture species.

### **Impacts of high stocking density on aquatic animals**

Stocking density is critical in determining a successful aquaculture practice (Bi et al., 2023). A high stocking density may be profitable and cost-effective in maximising space utilisation in an aquatic environment, but this practice forces the farmed species to compete for basic necessities such as food, space, and dissolved oxygen (Bi et al., 2023). Earlier studies claimed that a high stocking density could reduce fish growth performance and immunity, thus diminishing the total aquaculture production (Dediu et al., 2021). Therefore, stocking density has been recognised as a major source of chronic stress, which entails proper management for good aquaculture practices.

Stocking density may lead to biological or crowding stress, directly impacting fish gene expression, welfare, and production (Dediu et al., 2021; Jia et al., 2016). Furthermore, crowding stress disturbs the normal physiological response required for internal homeostasis maintenance (Bayunova et al., 2002; Long et al., 2019). Crowding stress also causes endocrine changes related to stress hormones, namely corticosteroids and catecholamines (Herrera et al.,

2019). Endocrine alterations can influence an organism's physiological aspects, such as haematological indices, antioxidant capacity, and immune system (Herrera et al., 2019), manifested as adverse effects on growth performance, disease resistance, behaviour, and survival (Bruce A. Barton, 2002) (see Figure 1).




Figure 1. Impacts of crowding stress on aquatic animals

### **Impacts of crowding stress on the growth performance of aquatic animals**

Crowding stress harms the growth performance of aquatic animals (see Table 1). Current research on the impact of crowding stress is limited to several aquatic species, such as sturgeon, *Acipenser queldenstaedtii*, Nile tilapia, *Oreochromis niloticus*, Senegalese sole, *Solea senegalensis*, Chinese sturgeon, *Acipenser sinensis*, hybrid sturgeon and minor carp, *Labeo bata*. The study outcomes indicated that high stocking density causes crowding stress, increased stress hormones, and glucose imbalance in aquatic animals, negatively impacting their growth performance. Nonetheless, there are several cases where crowding stress had no adverse effects on the farmed aquatic animals. These discrepancies warrant further investigation to substantiate the impact of crowding stress on the growth performance of other aquatic animal species, particularly crustaceans, due to the lack of literature.

At low stocking density, sturgeons (*A. queldenstaedtii*) demonstrated superior growth performance compared to their counterpart at high stocking density (Çelikkale et al., 2005). Despite the insignificant results, the low-stocking density group recorded a better final weight and survival rate than the high-density group. Likewise, studies on juvenile of Senegalese sole, *Solea senegalensis* (Andrade et al., 2015; Costas et al., 2013; Costas et al., 2008; Salas-Leiton et al., 2008), larvae and fingerling of lake sturgeon, *A. fulvescens* (Fajfer et al., 1999), Atlantic sturgeon, *A. oxyrinchus* (Mohler et al., 2000) reported similar findings. The insignificant results

in growth performance between the low and high stocking density groups may be attributed to the adaptation of the experimental fish to the high stocking density during the acclimatisation period. Another possible explanation is the fish farmed in high stocking density were fed with higher feed ration to fulfil their dietary requirements, thus sustaining themselves even under non-optimal conditions as observed in the Senegalese sole (Salas-Leiton et al., 2008). Similarly, (Costas et al., 2013) reported that Senegalese sole farms in high and low stocking exhibited comparable growth performance with the condition that the high stocking density fish are supplemented with a higher protein diet.

In some cases, high stocking density can promote the growth performance of aquatic animals, as reported in juvenile Siberian sturgeon, *A. baerii*. Despite that, high stocking density remains a risk for causing significantly ( $p < 0.05$ ) lower growth performance as observed in hybrid sturgeon farming (Bi et al., 2023; Dediu et al., 2021), Chinese sturgeon (Long et al., 2019), Atlantic cod (Lambert & Dutil, 2001), Nile tilapia (Gibtan et al., 2008), European seabass (Saillant et al., 2003) and rainbow trout (Ellis et al., 2002). In severe cases, crowding stress caused mortality in hybrid sturgeon (Bi et al., 2023), Nile tilapia, *Oreochromis niloticus* (Garcia et al., 2013), and Minor carp, *Labeo bata* (Karnatak et al., 2021). Thus, researchers proposed the use of the Biofloc system to ease crowding stress, which was proven effective in juvenile Common carp farmed at high stocking density (Adineh et al., 2022; Yousefi, Adineh, et al., 2023).

Glucose is an essential nutrient for animals that provides energy in the form of adenosine triphosphate (ATP). Many studies utilised the blood glucose level to indicate chronic stress, particularly crowding stress (IWAMA et al., 2015). High blood glucose levels have been linked to high stocking density (Long et al., 2019), but this is not the case for some studies as the species has acclimatised itself in a crowded environment (Dediu et al., 2021). Therefore, blood glucose levels in the blood may not be the most precise measurement of stress (Dediu et al., 2021).

Cortisol is an important hormone that regulates glucose production in aquatic animals via gluconeogenesis and glycogenolysis pathways (IWAMA et al., 2015). This stress hormone level can be used to reflect the stress experienced by an organism (H. Adineh et al., 2021; Fatima et al., 2017) and has been applied in earlier studies (Mommsen et al., 1999). An organism under stress will trigger the secretion of catecholamines hormone into the body (see Figure 2). Once present in blood, catecholamines stimulate the hypothalamic-pituitary-interrenal (HPI) to release cortisol into the circulatory system. Cortisol is responsible for mobilising energy reserves in the body by regulating and activating liver glycogenolysis and inhibiting glycolysis (Mazeaud et al., 1977). A high cortisol level will activate glycogenolysis and inhibit glycolysis, resulting in higher blood glucose production, which is common in aquatic animals under high stocking density (Dediu et al., 2021; Jia et al., 2016) and crowding stress (Ghafarifarsani et al., 2023; Hoseini, Yousefi, et al., 2019; Hosseini & Hoseini, 2012). These adaptive responses provide extra energy for cells in their bodies to handle crowding (B. A. Barton, 2002), hence reducing the growth performance of aquatic animals in high stocking density environments.

Haematological indices are also good stress indicators (Jeney & Jeney, 2002) for aquatic animals under high stocking density, including total haemoglobin, total red blood cells, high mean corpuscular haemoglobin concentration, and low mean corpuscular volume (Dediu

et al., 2021). The changes in haematological indices are adaptive responses for aquatic animals with high oxygen requirements and metabolic demands to maintain hemostasis (Dediu et al., 2021), particularly under crowding stress (Kleinert et al., 2019). Therefore, low-density aquatic animal farming remains ideal in aquaculture production, as Garcia et al. (2013) described in Nile tilapia, yielding superior production and uniform fish size. Nevertheless, several aquatic animals are adaptive to high-density farming, such as African catfish (Abdul Kari et al., 2021) and Senegalese sole (Andrade et al., 2015), with the condition that the farmed aquatic animals must be supplemented with high protein diet to mitigate crowding stress.




Figure 2. Mechanism of growth impairment of aquatic animals under crowding stress

Table 1. Impacts of crowding stress on the growth performance of aquatic animals

Species (stage)	Stocking density	Environment/ Duration	Impact on growth performance	References
Sturgeon, <i>Acipenser queldenstaedtii</i> (juvenile)	8 fish/m <sup>3</sup> (low density) & 12 fish /m <sup>3</sup> (high density)	Net cage (2m x 2m x 2m) 203 days	No significant difference between groups	(Çelikkale et al., 2005)
Nile tilapia, <i>Oreochromis niloticus</i> (juvenile)	50, 100, 150 & 200 fish /m <sup>3</sup>	Net cage 150 days	Low-density group demonstrated	(Gibtan et al., 2008)

			significantly better outcomes	
Nile tilapia, <i>Oreochromis niloticus</i> (juvenile)	Stage I: 800, 2000, 2500, 3000 fish/cage (6 m <sup>3</sup> )  Stage II: 80, 100, 120 kg/m <sup>3</sup>	Cage  Stage I: 41 days  Stage II: endpoint when the fish reached 800g weight	Low density showed significantly better results, and fish in uniform size  Disease prevalence was higher in the high stocking density group	(Garcia et al., 2013)
Senegalase sole, <i>Solea senegalensis</i> (juvenile)	Initial densities: 7, 17, and 24 kg/m <sup>2</sup>  Final densities: 13, 31, and 40 kg/m <sup>2</sup>	Raceway system  60 days	No significant difference between groups	(Andrade et al., 2015)
Chinese sturgeon, <i>Acipenser sinensis</i> (juvenile)	Low density: 4.80 kg/m <sup>2</sup>  Medium density: 8.99 kg/m <sup>2</sup>  High density: 12.68 kg/m <sup>2</sup>	Recirculating aquaculture system (RAS)  3 months	Low-density group exhibited significantly superior outcomes	(Long et al., 2019)
Hybrid sturgeon, (Bester ♀ x Beluga ♂ juvenile)	Trial 1: 2 & 4 kg/m <sup>2</sup>  Trial 2: 5 & 12 kg/m <sup>2</sup>	Recirculating aquaculture system (RAS)  6 weeks	Low-density group performed significantly better outcomes	(Dediu et al., 2021)
Minor carp, <i>Labeo bata</i> (juvenile)	50, 75, 100 fish/m <sup>3</sup>	Cage (5m x 5m x 3m)  180 days	Low-density group showed significantly better result	(Karnatak et al., 2021)

Hybrid sturgeon, ( <i>Acipenser baerii</i> ♀ X <i>Acipenser schrenckii</i> ♂ juvenile)	Low density: 100 - 150 fish/m <sup>3</sup>  Medium density: 200-250/m <sup>3</sup>  High density: 300 - 350/m <sup>3</sup>	In flow water pond  60 days	Low density group exhibited significantly better result	(Bi et al., 2023)
--	--	-----------------------------------	---	-------------------

---

### Impacts of crowding stress on the immune system and antioxidant capacity of aquatic animals

Aquatic animals farmed at high stocking density face the risk of immunosuppression and reduced antioxidant levels due to crowding stress (see Table 2), characterised by inhibiting metabolic and antioxidant enzyme activities. Despite evidence on the adverse effects of crowding stress on animal health and antioxidant status, several studies have reported no significant impacts on the farmed species. Therefore, further study is required to elucidate the effects of crowding stress on aquatic animals, particularly crustaceans.

Aquatic animals farmed in a high-density environment generally exhibit poorer immune function and a higher prevalence of disease infection (Aranguren et al., 2002; Mauri et al., 2011). For instance, Nile tilapia farmed in a high-density cage culture can exhibit higher disease infection and mortality (Garcia et al., 2013). High stocking density can also lead to immunosuppression, as observed in the common carp, *Cyprinus carpio* (H. Adineh et al., 2021; Hoseini, Abtahi, et al., 2019), turbot, *Scophthalmus maximus* (Jia et al., 2016), grass carp, *Ctenopharyngodon idella* (Li et al., 2023; Lin et al., 2018), and gilthead seabream, *Sparus aurata* (Montero et al., 1999). In addition, crowding stress causes diminishes red and white blood cells, leading to immune system impairment in Grass carp, *Ctenopharyngodon idella* (Lin et al., 2018). Nonetheless, immunosuppression did not occur in several studies on crowding stress, indicating the plasticity of the immune system that could be influenced by certain factors and metabolic changes (Al-Bakheit et al., 2016; Bi et al., 2023).

Crowding stress reportedly elevated serum biochemical parameters, such as glucose, cortisol, alkaline phosphatase (ALP), lactate dehydrogenase (LDH), alanine aminotransferase (ALT), and aspartate aminotransferase (AST) (Ghafarifarsani et al., 2023). Likewise, crowding stress elevated cortisol and glucose levels in common carp, *Cyprinus carpio* (Ghafarifarsani et al., 2023; Hoseini, Yousefi, et al., 2019; Hosseini & Hoseini, 2012) as an adaptive response and provide additional energy (B. A. Barton, 2002). A high cortisol level in fish leads to an impaired immune system (Tort, 2011). Crowding stress did not adversely impact the immune and antioxidant status of several aquatic animals, such as Senegalese sole, *Solea senegalensis*. (Andrade et al., 2015). There were no significant differences ( $p > 0.05$ ) between the high and low stocking density groups, and the hematocrit results demonstrated blood parameters such

as cortisol and glucose levels remained constant in both groups (Andrade et al., 2015). Similar cases were also recorded in other aquaculture species, including European seabass, *Dicentrarchus labrax* (Pascoli et al., 2011), gilthead seabream, *Sparus aurata* (Tort et al., 1998), and rainbow trout, *Oncorhynchus mykiss* (Morgan et al., 2008). These discrepancies suggest that aquatic animals are adaptive to a high stocking-density environment (Andrade et al., 2015).

Low lysozyme activity has been reported in aquatic animals under stressful environments with high stocking density or infected by disease, including Siberian sturgeon, *Acipenser baerii* (Chesneau, 2018) and Chinese sturgeon, *Acipenser sinensis* (Long et al., 2019) in high stocking density. Thus, lysozyme activity is a suitable indicator of the health status of aquatic animals. Moreover, a high stocking density could inhibit and block metabolic and antioxidant enzyme activities, inducing oxidative stress (Gao et al., 2017; Junqueira et al., 2004) and physiological imbalances in aquatic animals (E. Aksakal et al., 2011; Ercüment Aksakal et al., 2011). Despite that, several studies discovered no significant ( $p > 0.05$ ) difference in the antioxidant capacity of tongue sole, *Cynoglossus semilaevis* (Liu et al., 2016), largemouth bass, *Micropterus salmonides*, and Senegalese sole, *Solea senegalensis* (Andrade et al., 2015) at different stocking densities, suggesting the adaptability of farmed fish under high stocking density.

High stocking density also reportedly triggers oxidative stress and induces lipid peroxidation activity. For instance, antioxidant capacity induced lipid peroxidation and reduced superoxide dismutase (SOD) and glutathione (GSH-PX) activities in the serum of hybrid sturgeon farmed under high stocking density (Bi et al., 2023; Liu et al., 2014; Long et al., 2019; Yousefi et al., 2020). Lipid peroxidation produces malondialdehyde (MDA), a metabolite that can be measured in blood serum to indicate oxidative stress level. In addition, downregulation of skin immune-related genes, such as interleukin - 1 $\beta$  (IL-1 $\beta$ ), insulin -like growth factor – (IGF-), and lysozyme (LZM), were observed in juvenile Turbot, *Scophthalmus maximus* under high stocking density (Jia et al., 2016). The study also found that high stocking density inhibits mucus enzyme activities, induces oxidative stress in the skin, and decreases lysozyme activity in juvenile Turbot (Jia et al., 2016).

Heat shock protein (Hsp) expression parallels stress exposure; hence, it is a good stress indicator for aquatic animals (Sanders, 1993). The Hsps are classified based on molecular weight: small Hsps (< 40kDa), Hsp60, Hsp70, Hsp90, and Hsp100. These proteins are produced as a protective mechanism against protein denaturation in the events of hyperthermia, cytotoxicity from pesticides, heavy metals, ethanol, and hypoxia (Dorts et al., 2009; Lee & Corry, 1998; Lewis et al., 1999; Lindquist & Craig, 1988), and crowding stress (Ercüment Aksakal et al., 2011).

Table 2. Impacts of crowding stress on the immune system and antioxidant capacity of aquatic animals

Species (stage)	Stocking density	Environment/ Duration	Impacts on the immune system and antioxidant capacity	References
-----------------	------------------	--------------------------	---	------------

Rainbow trout, <i>Oncorhynchus mykiss</i> (juvenile)	Low stocking density: 10 kg/m <sup>3</sup>	9 months	Blood cortisol level increases in the high stocking density group	(North et al., 2006)
	High stocking density: 40 and 80 kg/m <sup>3</sup>		Lower lysozyme activity in the high stocking density group	
Rainbow trout (juvenile)	Control group: 15 kg/m <sup>3</sup>	Fiber glass tank 1m (diameter) x 1.2m (depth)	High stocking density inhibits metabolic and antioxidant enzyme activities	(Ercüment Aksakal et al., 2011)
	High stocking density: 20, 25, and 30 kg/m <sup>3</sup>	2 months	High stocking density upregulates heat shock protein (Hsp70) expression	
Senegalase sole, <i>Solea senegalensis</i> (juvenile)	Initial densities: 7, 17, and 24 kg/m <sup>2</sup>	Raceway system  60 days	No significant difference between groups	(Andrade et al., 2015)
	Final densities: 13, 31, and 40 kg/m <sup>2</sup>			
Turbot, <i>Scophthalmus maximus</i> (juvenile)	Low density: 9.3 kg/m <sup>2</sup>	Recirculating aquaculture system (RAS)	High stocking density inhibits mucus enzyme activities	(Jia et al., 2016)
	Medium density: 13.6 kg/m <sup>2</sup>	120 days	High stocking density induces oxidative stress on the skin	

	High density: 19.1 kg/m <sup>2</sup>		High stocking density downregulated the expression of skin immune-related gene expression	
			High stocking density decreases lysozyme activity	
Grass carp, <i>Ctenopharyngodon idella</i> (juvenile, 98.48g)	Low density: 0.9 kg/m <sup>2</sup>	Tank 10 weeks	High stocking density induces immunosuppression	(Lin et al., 2018)
	Medium density: – 2.97 kg/m <sup>2</sup>			
	High density: 5.9 kg/m <sup>2</sup>			
Chinese sturgeon, <i>Acipenser sinensis</i> (juvenile)	Low density: 4.80 kg/m <sup>2</sup>	RAS 3 months	High stocking density decreases lysozyme activity	(Long et al., 2019)
	Medium density: 8.99 kg/m <sup>2</sup>			
	High density: 12.68 kg/m <sup>2</sup>			

Hybrid sturgeon, ( <i>Acipenser baerii</i> ♀ X <i>Acipenser</i> <i>schrenckii</i> ♂ juvenile)	Low density: 100 - 150 fish/ m <sup>3</sup>	In-flow water pond 60 days	No significant difference in immune status	(Bi et al., 2023)
	Medium density: 200 – 250 fish/ m <sup>3</sup>		High stocking density inhibits antioxidant capacity	
	High density: 300 – 350 fish/m <sup>3</sup>			
Grass carp, <i>Ctenopharyngodon</i> <i>Idella</i> (712.70g)	Low density: 10kg/ m <sup>3</sup>	120 days	High stocking density induces immunosuppression	(Li et al., 2023)
	Medium density: 15kg/ m <sup>3</sup>			
	High density: 20kg/ m <sup>3</sup>			

### **Roles of phytobiotics in mitigating crowding stress among aquatic animals**

Various plant extracts can be used as phytobiotics to mitigate crowding stress in aquaculture (see Table 3). Phytobiotics can be utilised as a feed additive between 40 and 112 days to enhance the immune system and antioxidant capacity of the aquatic species. Furthermore, phytobiotics could improve blood parameters, immune system, antioxidant capacity, and disease resistance and mitigate crowding stress in aquaculture. Nevertheless, studies in this area are still in their infancy and require further investigation to explore the potential of phytobiotics in aquaculture.

Earlier studies have reported that Oak, *Quercus castaneifolia* leaf extract has high antioxidant and antibacterial properties (Almeida et al., 2008; Bahador & Baserisalehi, 2011) that could reduce free radical formation and inhibit the growth of pathogenic bacteria. Besides improving growth performance, the plant extract enhanced the antioxidant enzyme activities

such as SOD, CAT, and GPx to mitigate oxidative stress in the common carp (Paray et al., 2020). Oak leaf extract may also stimulate immune components, such as lysozyme elevation, to increase disease resistance in the fish (Hoseinifar et al., 2019; Paray et al., 2020). Conversely, oak leaf extract could not regulate the cortisol level of fish under stress for faster recovery compared to the untreated control group (Paray et al., 2020). Meanwhile, dietary *Rheum officinale* extract significantly regulated cortisol and glucose levels to mitigate crowding stress in common carp (J. Xie et al., 2008). Green tea extract also mitigated anaesthesia-induced stress in black rockfish, *Sebastes schlegeli* (Hwang et al., 2013), while *Urtica dioica* extract significantly reduced crowding stress in Victoria labeo, *Labeo victorianus* by lowering cortisol and glucose levels (Ngugi et al., 2015).

*Rheum officinale* bail is a Chinese herbal medicine for constipation, jaundice, ascites, and asthma (Akbar, 2020). Previous studies have claimed this herbal remedy enhances the immune system and relieves stress (Jian & Wu, 2004; Yin et al., 2006). Anthraquinone extract from rhubarb bail potentially improves blood parameters, disease resistance against *Aeromonas hydrophila*, and crowding stress in aquaculture (Jun Xie et al., 2008). Nevertheless, overusing Chinese herbs in farmed aquatic species causes gut microbiota imbalance (Liu et al., 2004) and affects the palatability of medicated feed, reducing their growth performance (Jun Xie et al., 2008).

Lemon (*Citrus limon*) peel reportedly possesses medicinal value such as antimicrobial, anticancer, and antioxidant properties (Espina et al., 2011). This high-fibre agricultural waste has been widely explored as a feed additive in aquaculture. Previous studies revealed that dried lemon peel (DLP) positively impacted the growth performance, haematological parameters, and immune system in various aquatic animals, such as common carp, *Cyprinus carpio* (Sadeghi et al., 2021), Nile tilapia, *Oreochromis niloticus*, African catfish, *Clarias gariepinus* (Abdel Rahman et al., 2019), gilthead seabream, *Sparus aurata* (García Beltrán et al., 2017) and Rohu, *Labeo rohita* (Harikrishnan et al., 2020).

DLP potentially mitigated crowding stress in rainbow trout, *Oncorhynchus mykiss*, by enhancing their immune system and antioxidant capacity, besides lowering the blood cortisol level (Chekani et al., 2021). The cortisol level is significantly lower in the group of fish that received DLP than the fish from the control group, which was also observed in juvenile *Labeo victorianus* (Ngugi et al., 2017). On the other hand, there was no significant ( $p > 0.05$ ) difference in cortisol levels between the DLP-treated and untreated Nile tilapia and African catfish (Abdel Rahman et al., 2019). Dietary DLP also significantly ( $P < 0.05$ ) increased the total immunoglobulin, alternative complement (ACH50), and lysozyme activity in rainbow trout than the control group. Currently, the mechanism of DLP in boosting the immune system remains clear, but studies have explained that these enhancements are caused by the bioactive compounds found in DLP (Baba et al., 2016; García Beltrán et al., 2017; Xi et al., 2017). The DLP also enhances the antioxidant capacity of aquaculture species by significantly increasing the SOD and CAT activities ( $p < 0.05$ ) compared to the control group. The SOD and CAT are the frontliners of the antioxidant defence against pathogenic stress (Zhang et al., 2013).

Rosemary, *Rosmarinus officinalis*, leaf is another phyto-biotic that can be used as a feed additive to reduce crowding stress in common carp due to the high stocking density farming system. This medicinal herb contains bioactive compounds with various health benefits, including flavonoids and tannins, which act as free radical scavengers and inhibit lipid

peroxidation (Almeida et al., 2009; Benedi et al., 2004; Zheleva-Dimitrova et al., 2010). For instance, dietary rosemary leaf demonstrated the ability to control cortisol and glucose levels in fish under crowding stress (Yousefi et al., 2019). Furthermore, combining Saint John wort, *Hypericum perforatum*, lemon balm, *Melissa officinalis*, and rosemary promoted the antioxidant capacity and mitigated crowding stress in juvenile Atlantic salmon, *Salmo salar*. Phenolic compounds, particularly in rosemary and lemon balm, contributed to antioxidant capacity enhancement in fish (Barros et al., 2013; Peng et al., 2007).

Pomegranate peel is another promising feed additive to ease crowding stress. This plant possesses various bioactive compounds, such as polyphenols, which are known for their pharmacological effects, including anti-diabetic, antioxidant and immunostimulatory (Akhtar et al., 2015; Malviya et al., 2014; Singh et al., 2019). Furthermore, studies have revealed that dietary pomegranate peel positively impacted the growth and health of Nile tilapia and rainbow trout (Avazeh et al., 2021; Badawi & Gomaa, 2016; Monir et al., 2020). This agriculture by-product also effectively lowered the plasma cortisol glucose in the blood of common carp subjected to crowding stress (Yousefi, Hoseini, et al., 2023).

Table 3. Roles of phytobiotics in mitigating crowding stress among aquatic animals

Phytobiotics/ (dosage)	Application/ Duration	Aquatic animals – Stage	Impacts	References
Anthraquinone extract from rhubarb <i>Rheum officinale</i> bail (1% and 2%)	Feed additive  70 days	Common carp, <i>Cyprinus carpio</i> var Jian  Juvenile (5.39g)	Improve blood parameters  Stimulate disease resistance against <i>Aeromonas hydrophila</i>  Mitigate crowding stress	(Jun Xie et al., 2008)
Stinging nettle, <i>Urtica dioica</i> (1%, 2%, and 5% of diet)	Feed additive  16 weeks	Victoria labeo, <i>Labeo victorianus</i>  Juvenile (25g)  Female adult (3.4kg)	Improve blood parameters  Stimulate disease resistance against <i>A. hydrophila</i>	(Ngugi et al., 2015)

		Male adult (470g)		
Saint John wort, <i>Hypericum perforatum</i> , lemon balm, <i>Melissa officinalis</i> , rosemary, <i>Rosmarinus officinalis</i> (ratio 3:2:1, 6g/kg of feed)	Feed additive  40 days	Atlantic salmon, <i>Salmo salar</i>  Juvenile (4.83g)	Enhance antioxidant capacity  Mitigate crowding stress	(Reyes-Cerpa et al., 2018)
Rosemary, <i>Rosmarinus officinalis</i> leaf (1-3% of feed)	Feed additive  65 days	Common carp, <i>Cyprinus carpio</i>  Juvenile (12.93g)	Improve growth, enhance immune system, increase antioxidant capacity, mitigate crowding stress	(Yousefi et al., 2019)
Oak, <i>Quercus castaneifolia</i> leaf extract (0.5, 1, and 2g/kg of feed)	Feed additive  60 days	Common carp, <i>Cyprinus carpio</i>  Juvenile (20g)	Enhance antioxidant and immune system  No significant improvement in growth performance  Mitigate crowding stress	(Paray et al., 2020)
Dehydrated <i>Citrus lemon</i> peel (DLP) (0.5 - 1.5% of diet)	Feed additive  45 days	Rainbow trout, <i>Oncorhynchus mykiss</i>  Juvenile (26.8g)	No significant improvement in growth performance	(Chekani et al., 2021)

			Enhance antioxidant and immune system	
			Improve blood parameters	
			Mitigate crowding stress	
Pomegranate peel (0.5% of diet)	Feed additive 56 days	Common carp, <i>Cyprinus carpio</i>	No adverse impact on the growth performance	(Yousefi, Hoseini, et al., 2023)
		Juvenile (18.1g)	Enhance antioxidant and immune system	
			Improve blood parameters	
			Mitigate crowding stress	

### Roles of probiotics in mitigating crowding stress in aquatic animals

*Shewanella putrefaciens* Pdp11 significantly ( $p < 0.05$ ) mitigated crowding stress by upregulating immune gene transcription, modulating gut microbiota, and stimulating disease resistance against *Vibrio harveyi* and *V. parahaemolyticus* in *Solea senegalensis* (Tapia-Paniagua et al., 2014) (see Table 4). Dietary *S. putrefaciens* Pdp11 also upregulated immune-related genes, such as C7, HP, NARS, HAMP1, TNFAIP9, and NCCRP1, responsible for producing antimicrobial peptides against pathogenic invasion. Likewise, these genes were upregulated in *S. senegalensis* lipopolysaccharide supplementation (Osuna-Jiménez et al., 2009; Prieto-Álamo et al., 2009). Meanwhile, *S. putrefaciens* Pdp11 potentially reduced the harmful bacteria family *Actinobacteria*, *Photobacterium*, and *Vibrio* spp. in the intestines of *S. senegalensis* (García de La Banda et al., 2010; Tapia-Paniagua et al., 2010).

Table 4. Roles of probiotics in mitigating crowding stress among aquatic animals

Probiotics/prebiotic s - Dosage	Application/Duratio n	Aquatic animals/	Impacts	Reference s
------------------------------------	--------------------------	---------------------	---------	----------------

		stages (weight)		
<i>Shewanella putrefaciens</i> Pdp11 – 10 <sup>9</sup> CFU/g	Feed additive 30 days	<i>Solea senegalensis</i> Juvenile (14.6g)	Mitigate crowding stress Upregulate immune gene transcription Modulate gut microbiota Stimulate disease resistance against <i>Vibrio harveyi</i> and <i>V. parahaemolyticus</i>	(Tapia-Paniagua et al., 2014)

### **Roles of synthetic immune stimulants (SIS) in mitigating crowding stress in aquatic animals**

Dietary gallic acid significantly reduced glucose and cortisol levels in common carp, *Cyprinus carpio*, to mitigate crowding stress (Ghafarifarsani et al., 2023) (see Table 5). A similar finding has been reported in broilers (Biswas et al., 2023). However, the mechanism of gallic acid in relieving crowding stress remains unclear. Dietary yeast, *Xanthophyllomyces dendrorhous* (sexual stage *Phaffia rhodozyma*) extract at a dose of 0.5g/kg of feed for 40 days enhanced the antioxidant capacity and mitigate crowding stress in juvenile Atlantic salmon, *Salmo salar*.

Yeast extract contains bioactive compounds, primarily carotenoids. Carotenoids are vitamin A precursors acting as free radical scavengers to eliminate excess energy, preventing cellular oxidative damage (Amengual et al., 2011; Loto et al., 2012; Martinez-Moya et al., 2015). Astaxanthin is the main bioactive compound in yeast extract, a carotenoid that can inhibit lipid peroxidation activity (Contreras et al., 2015). Other carotenoids in yeast extract include lycopene,  $\beta$ -carotene, lycopene, and phoenicoxanthin (Contreras et al., 2015; Córdova et al., 2016). The  $\beta$ -glucan is a primary bioactive compound in the yeast cell wall, known for the antioxidant properties and MDA inhibition (Lee et al., 2011). Therefore, astaxanthin and  $\beta$ -glucan in the yeast extract may be responsible for boosting the antioxidant capacity of the experimental fish by inhibiting lipid peroxidation, lowering MDA level, protecting cells from oxidative damage, and mitigating crowding stress.

The supplementation of dietary pennyroyal, *Mentha pulegium*, essential oil improved the growth performance, enhanced the immune system, increased antioxidant capacity, and mitigated crowding stress in common carp (Yousefi, Adineh, et al., 2023). Pennyroyal essential

oil was found to activate digestive enzyme activity in the fish and promote gut health, enhancing the fish's growth performance (Kaur et al., 2018; Ringø et al., 2016). In addition, pulegone, an organic compound that is abundant in essential oil, was reported to enhance antioxidative response and ease stress due to crowding in the fish (Yousefi, Adineh, et al., 2023).

Chrysin is a bioactive compound present in various natural sources such as honey, propolis, blue passion, and mushrooms (Mani & Natesan, 2018). The medicinal properties of this compound have been widely reported, including immunostimulatory, anti-inflammatory, antimicrobial, and antioxidant activities (Stompor-gorący et al., 2021; Zeinali et al., 2017). Thus, chrysin could be included as a feed additive to boost fish production and mitigate crowding stress in an aquaculture system. In a previous study, rainbow trout that received a diet containing chrysin for 56 days demonstrated improved antioxidative response and immune system (Yousefi, Nedaei, et al., 2023).

Table 5. Roles of synthetic immune stimulants (SIS) in mitigating crowding stress in aquatic animals

SIS - Dosage	Application/Duration	Aquatic animals/ stage	Impacts	References
Gallic acid – 150, 300, 450 & 600 mg/kg of feed	Feed additive – 60 days	Common carp, <i>Cyprinus carpio</i> - juvenile	Improve growth, enhance immune system, increase antioxidant capacity, mitigate crowding stress	(Ghafariarsani et al., 2023)
Yeast, <i>Xanthophyllomyces dendrorhous</i> (sexual stage <i>Phaffia rhodozyma</i> ) extract – 0.5g/kg of feed	Feed additive – 40 days	Atlantic salmon, <i>Salmo salar</i> , juvenile (4.83g)	Enhance antioxidant capacity  Mitigate crowding stress	(Reyes-Cerpa et al., 2018)
Pennyroyal essential oil - 250mg/kg of feed	Feed additive – 56 days	Common carp, <i>Cyprinus carpio</i> - juvenile (13g)	Improve growth, enhance immune	(Yousefi, Adineh, et al., 2023)

system,  
increase  
antioxidant  
capacity,  
mitigate  
crowding  
stress

Chrysin – 100 -  
700mg/kg of feed

Feed additive – 56  
days

Rainbow  
trout,  
*Oncorhynchus*  
*mykiss*

Juvenile (51g)

Improve  
growth,  
enhance  
immune  
system,  
increase  
antioxidant  
capacity,  
mitigate  
crowding  
stress

(Yousefi,  
Nedaei, et al.,  
2023)

---

## Roles of feed additives combination in mitigating crowding stress in aquatic animals

A probiotic combination of *Lactobacillus acidophilus* (LAB) and *resveratrol* (RE) was found to mitigate crowding stress in common carp, *Cyprinus carpio* (Gabr et al., 2023) (Table 6). The functional roles of LAB include enhancing digestion and nutrient absorption, digestive enzyme activity, growth-related metabolite production, and eliminating harmful gut bacteria (Assan et al., 2022; Balcázar et al., 2006). Meanwhile, RE can ameliorate intestinal damage, reduce protein degradation, enhance antioxidant capacity, and improve lipid and glucose metabolism (Zhang et al., 2018) (Salomão et al., 2019; Tan et al., 2019; Wilson et al., 2015). Furthermore, the LAB and RE combination enhanced the antioxidant capacity by increasing antioxidant enzymes to protect the antioxidant defence system, thus relieving crowding stress in common carp (Gabr et al., 2023). Previous studies also revealed that probiotics can protect the antioxidant mechanism against free radicals and oxidative stress in fish (Hoseinifar et al., 2020) and other animals (Heshmati et al., 2018).

Combining selenium and garlic extract as feed additives positively impacted Grass carp, *Ctenopharyngodon Idella*, exhibiting improved growth performance, antioxidative response, and mitigated crowding stress (Hossein Adineh et al., 2021). Selenium was reported as an essential component of glutathione peroxide enzyme, which protects the cellular membrane from oxidative impairment (Rotruck et al., 1973). Meanwhile, garlic is a widely used phytobiotic that is utilised as a feed additive to enhance the growth and health of various aquatic species (Hossein Adineh et al., 2021). Flavonoids and sulphur are two primary components of garlic that are rich in antioxidants and can eliminate free radicals in fish (Chung, 2006; Sharma et al., 2010). Therefore, the selenium and garlic combination is ideal in fish farming.

Table 6. Roles of feed additives combination in mitigating crowding stress in aquatic animals

Feed additives - Dosage	Application/Duration	Aquatic animals – stages (weight)	Impacts	References
<i>Lactobacillus acidophilus</i> (LAB) + resveratrol (RE) - 1.5 X 10 <sup>7</sup> CFU/g + 300mg/kg	Feed additive 60 days	Common carp, <i>Cyprinus carpio</i> Juvenile (30.16g)	Mitigate crowding stress	(Gabr et al., 2023)
Nano selenium + garlic extract - 1-2 mg + 1-2 g	Feed additive 60 days	Grass carp, <i>Ctenopharyngodon idella</i> Juvenile (2.1g)	Improve growth, increase antioxidant capacity,	(Hossein Adineh et al., 2021)

### **Conclusion**

High stocking density in aquaculture causes crowding stress in farmed species, impairing their growth performance and health status. Researchers have proposed using feed additives, namely probiotics, phytobiotics, and SIS, to mitigate this issue by modulating the immune system, antioxidant capacity, and intestinal condition of aquatic animals. Furthermore, feed additives have exhibited protective effects against crowding stress, but the mechanism of these components has yet to be fully elucidated. Moreover, limited studies have explored the use of probiotics as feed additives to mitigate crowding stress in fish farming.

Currently, studies have revealed that feed additives can potentially control cortisol and glucose levels in fish after being exposed to crowding stress. Most studies incorporated phytobiotic as feed additives between 0.5% and 5% of the diet to manage the effects of crowding stress in fish farming. Alternatively, SIS has demonstrated the potential as a feed additive and reducing stress due to crowding in fish farming at an even lower inclusion level ( $\leq 0.5\%$  of diet). A combination of SIS and probiotics or phytobiotics also demonstrated positive outcomes in mitigating stress in fish farming. Future studies should investigate the appropriate dosage and optimise feed additives administration to identify the best practices for sustainable aquaculture production. Furthermore, future studies should continue to explore the application of the Biofloc technology in fish farming, which was found to reduce crowding stress due to high stocking density.

### **Acknowledgement**

This research was funded by the Matching Grant Universiti Malaysia Kelantan (R/MTCH/A0700/00387A/009/2023/01161) and the Ministry of Higher Education, Malaysia, under the Niche Research Grant Scheme (NRGS) (R/NRGS/A0.700/00387A/006/2014/00152).

### **Author contributions**

All authors contributed to the study conception and design. Lee Seong Wei, Martina Irwan Khoo, Ramasamy Harikrishnan, Ümit Acar, Md Eilious Hosain, Azra MN, Ahmad Syazni Kamarudin, Lau Kia Kian, and Wendy Wee performed material preparation, data collection, and analysis. Lee Seong Wei and Wendy Wee wrote the first draft of the manuscript, while Martina Irwan Khoo commented on previous versions. All authors have read and approved the final manuscript.

### **Conflicts of interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- Abdel Rahman A.N., ElHady M., Shalaby S.I. (2019). Efficacy of the dehydrated lemon peels on the immunity, enzymatic antioxidant capacity and growth of Nile tilapia (*Oreochromis niloticus*) and African catfish (*Clarias gariepinus*). *Aquaculture*, 505: 92–97.
- Abdul Kari Z., Kabir M.A., Mat K., Rusli N.D., Razab M.K.A.A., Ariff N.S.N.A., Edinur H.A., Rahim M.Z.A., Pati S., Dawood M.A.O., Wei L.S. (2021). The possibility of replacing fish meal with fermented soy pulp on the growth performance, blood biochemistry, liver, and intestinal morphology of African catfish (*Clarias gariepinus*). *Aquacult. Rep.*, 21: 100815.
- Abdul Kari Z., Wee W., Mohamad Sukri S.A., Che Harun H., Hanif Reduan M.F., Irwan Khoo M., Van Doan H., Wen Goh K., Seong Wei L. (2022). Role of phytobiotics in relieving the impacts of *Aeromonas hydrophila* infection on aquatic animals: A mini-review. *Front. Vet. Sci.*, 9.
- Adineh H., Naderi M., Jafaryan H., Khademi Hamidi M., Yousefi M., Ahmadifar E. (2022). Effect of Stocking Density and Dietary Protein Level in Biofloc System on the Growth, Digestive and Antioxidant Enzyme Activities, Health, and Resistance to Acute Crowding Stress in Juvenile Common Carp (*Cyprinus carpio*). *Aquacul. Nutr.*, 9344478.
- Adineh H., Naderi M., Nazer A., Yousefi M., Ahmadifar E. (2021). Interactive effects of stocking density and dietary supplementation with Nano selenium and garlic extract on growth, feed utilisation, digestive enzymes, stress responses, and antioxidant capacity of grass carp. *J. World Aquacul. Soc.*, 52: 789–804.
- Adineh H., Naderi M., Yousefi M., Khademi Hamidi M., Ahmadifar E., Hoseini S.M. (2021). Dietary licorice (*Glycyrrhiza glabra*) improves growth, lipid metabolism, antioxidant and immune responses, and resistance to crowding stress in common carp, *Cyprinus carpio*. *Aquacul. Nutr.*, 27: 417–426.
- Akbar S. (2020). *Rheum officinale* Baill.; *R. palmatum* L. (*Polygonaceae*). In S. Akbar (Ed.), *Handbook of 200 Medicinal Plants: A Comprehensive Review of Their Traditional Medical Uses and Scientific Justifications* (pp. 1527–1537). Springer International Publishing.
- Akhtar S., Ismail T., Fraternali D., Sestili P. (2015). Pomegranate peel and peel extracts: Chemistry and food features. *Food Chem.*, 174: 417–425.
- Aksakal E., Ekinci D., Erdogan O., Beydemir S., Alim Z., Ceyhun S.B. (2011). Increasing stocking density causes inhibition of metabolic-antioxidant enzymes and elevates mRNA levels of heat shock protein 70 in rainbow trout. *Livestock Sci.*, 141: 69–75.
- Al-Bakheit A.A., Traka M., Saha S., Mithen R., Melchini, A. (2016). Accumulation of Palmitoylcarnitine and Its Effect on Pro-Inflammatory Pathways and Calcium Influx in Prostate Cancer. *Prostate.*, 76: 1326–1337.

- Almeida I.F., Fernandes E., Lima J.L.F.C., Costa P.C., Bahia M.F. (2008). Protective effect of *Castanea sativa* and *Quercus robur* leaf extracts against oxygen and nitrogen reactive species. *J. Photochem. Photobiol. B: Biol.*, 91: 87–95.
- Amano M., Amiya N., Fukushima K., Hagio H., Yamamoto N., Sakakura Y. (2022). Effects of crowding stress on the hypothalamo-pituitary-interrenal axis of the self-fertilising fish, *Kryptolebias marmoratus*. *Comp. Biochem. Physiol. Part A: Molec. Integra. Physiol.*, 264: 111110.
- Amengual J., Lobo G.P., Golczak M., Li H.N.M., Klimova T., Hoppel C.L., Wyss A., Palczewski K., Von Lintig J. (2011). A mitochondrial enzyme degrades Carotenoids and protects against oxidative stress. *FASEB*, 25: 948–959.
- Andrade T., Afonso A., Pérez-Jiménez A., Oliva-Teles, A., de las Heras V., Mancera J.M., Serradeiro R., Costas B. (2015). Evaluation of different stocking densities in a Senegalese sole (*Solea senegalensis*) farm: Implications for growth, humoral immune parameters and oxidative status. *Aquaculture*, 438: 6–11.
- Anis Mohamad Sukri S., Andu Y., Tuan Harith Z., Sarijan S., Naim Firdaus Pauzi M., Seong Wei L., Dawood M.A.O., Abdul Kari Z. (2022). Effect of feeding pineapple waste on growth performance, texture quality and flesh colour of Nile tilapia (*Oreochromis niloticus*) fingerlings. *Saudi J. Biol. Sci.*, 29: 2514–2519.
- Aranguren R., Tafalla C., Novoa B., Figueras A. (2002). Experimental transmission of encephalopathy and retinopathy induced by nodavirus to sea bream, *Sparus aurata* L., using different infection models. *J. Fish Dis.*, 25: 317–324.
- Assan D., Kuebutornye F.K.A., Hlorodzi V., Chen H., Mraz J., Mustapha U.F., Abarike E.D. (2022). Effects of probiotics on digestive enzymes of fish (finfish and shellfish); status and prospects: a mini review. *Comp. Biochem. Physiol. Part - B: Biochem. Molec. Biol.*, 257: 110653.
- Avazeh A., Adel M., Shekarabi S.P.H., Emamadi H., Dawood M.A.O., Omidi A.H., Bavarsad M. (2021). Effects of dietary pomegranate peel meal on the growth performance, blood indices, and innate immune response of rainbow trout (*Oncorhynchus mykiss*). *Ann. Anim. Sci.*, 21: 233–244.
- Baba E., Acar Ü., Öntaş C., Kesbiç O. S., Yılmaz S. (2016). Evaluation of *Citrus limon* peels essential oil on growth performance, immune response of Mozambique tilapia *Oreochromis mossambicus* challenged with *Edwardsiella tarda*. *Aquaculture*, 465: 13–18.
- Badawi M.E., Gomaa A.M. (2016). Influence of diets supplemented with pomegranate peel extract on performance in *Oreochromis niloticus*. *Japanese J. Vet. Res*, 64, S87–S94.
- Bahador N., Baserisalehi M. (2011). The effect of *Quercus castaneifolia* extract on pathogenic enteric bacteria. *Anaerobe*, 17: 358–360.
- Balcázar J. L., Vendrell D., de Blas I., Ruiz-Zarzuola I., Gironés O., Muzquiz J.L. (2006). Immune modulation by probiotic strains: Quantification of phagocytosis of *Aeromonas salmonicida* by leukocytes isolated from gut of rainbow trout (*Oncorhynchus mykiss*) using a radiolabelling assay. *Comp. Immunol., Microbiol. Infect. Dis.*, 29: 335–343.
- Barros L., Dueñas M., Dias M.I., Sousa M.J., Santos-Buelga C., Ferreira I.C.F.R. (2013). Phenolic profiles of cultivated, *in vitro* cultured and commercial samples of *Melissa officinalis* L. infusions. *Food Chem.*, 136: 1–8.

- Barton B.A. (2002). Stress in fishes: A diversity of responses with particular reference to changes in circulating corticosteroids. *Integr. Comp. Biol.*, 42: 517–525.
- Bayunova L., Barannikova I., Semenkova T. (2002). Sturgeon stress reactions in aquaculture. *J. Appl. Ichthyol.*, 18: 397–404.
- Benedí J., Arroyo R., Romero C., Martín-Aragón S., Villar A.M. (2004). Antioxidant properties and protective effects of a standardised extract of *Hypericum perforatum* on hydrogen peroxide-induced oxidative damage in PC12 cells. *Life Sci.*, 75: 1263–1276.
- Bi B., Yuan Y., Zhao Y., He M., Song H., Kong L., Gao Y. (2023). Effect of crowding stress on growth performance, the antioxidant system and humoral immunity in hybrid sturgeon. *Aquacul. Rep.*, 28: 101468.
- Biswas S., Cho S.B., Kim I. H. (2023). An evaluation of gallic acid supplementation to corn-soybean-gluten meal-based diet in broilers. *Poultry Sci.*, 102: 102738.
- Castro-Osses D., Carrera-Naipil C., Gallardo-Escárate C., Gonçalves A.T. (2017). Functional diets modulate the acute phase protein response in *Oncorhynchus mykiss* subjected to chronic stress and challenged with *Vibrio anguillarum*. *Fish. Shellfish. Immunol.*, 66: 62–70.
- Çelikkale M.S., Memiş D., Ercan E., Çağıltay F. (2005). Growth performance of juvenile Russian sturgeon (*Acipenser gueldenstaedtii* Brandt & Ratzenburg, 1833) at two stocking densities in net cages. *J. Appl. Ichthyol.*, 21: 14–18.
- Chekani R., Akrami R., Ghiasvand Z., Chitsaz H., Jorjani S. (2021). Effect of dietary dehydrated lemon peel (*Citrus limon*) supplementation on growth, hemato-immunological and antioxidant status of rainbow trout (*Oncorhynchus mykiss*) under exposure to crowding stress. *Aquaculture*, 539: 736597.
- Chen R., Lochmann R., Goodwin A., Praveen K., Dabrowski K., Lee K.J. (2004). Effects of dietary vitamins C and E on alternative complement activity, hematology, tissue composition, vitamin concentrations and response to heat stress in juvenile golden shiner (*Notemigonus crysoleucas*). *Aquaculture*, 242: 553–569.
- Chen X.M., Guo G.L., Sun L., Yang Q.S., Wang G.Q., Qin G.X., Zhang D.M. (2016). Effects of Ala-Gln feeding strategies on growth, metabolism, and crowding stress resistance of juvenile *Cyprinus carpio* var. Jian. *Fish. Shellfish. Immunol.*, 51: 365–372.
- Chesneau V. (2018). Immunology in Sturgeons with a Focus on the Siberian Sturgeon Mechanisms, Responses to Stress and Stimulation. In P. Williot, G. Nonnotte, & M. Chebanov (Eds.), *The Siberian Sturgeon (Acipenser baerii, Brandt, 1869) Volume 2 - Farming* (pp. 373–402). Springer International Publishing.
- Chung L.Y. (2006). The antioxidant properties of garlic compounds: allyl cysteine, alliin, allicin, and allyl disulfide. *J. Med. Food.*, 9: 205–213.
- Contreras G., Barahona S., Sepúlveda D., Baeza M., Cifuentes V., Alcaíno J. (2015). Identification and analysis of metabolite production with biotechnological potential in *Xanthophyllomyces dendrorhous* isolates. *World J. Microbiol. Biotechnol.*, 31: 517–526.
- Córdova P., Alcaíno J., Bravo N., Barahona S., Sepúlveda D., Fernández-Lobato M., Baeza M., Cifuentes V. (2016). Regulation of carotenogenesis in the red yeast *Xanthophyllomyces dendrorhous*: The role of the transcriptional co-repressor complex Cyc8-Tup1 involved in catabolic repression. *Microbial. Cell. Fact.*, 15: 193.

- Costas B., Aragão C., Dias J., Afonso A., Conceição L.E.C. (2013). Interactive effects of a high-quality protein diet and high stocking density on the stress response and some innate immune parameters of Senegalese sole *Solea senegalensis*. *Fish Physiol. Biochem.*, 39: 1141–1151.
- Costas B., Aragão C., Mancera J.M., Dinis M.T., Conceição L.E.C., Refojos, B.C. (2008). High stocking density induces crowding stress and affects amino acid metabolism in Senegalese sole *Solea senegalensis* (Kaup 1858) juveniles. *Aquacult. Res.*, 39: 1–9.
- Costas B., Aragão C., Soengas J.L., Míguez J.M., Rema P., Dias J., Afonso A., Conceição L.E.C. (2012). Effects of dietary amino acids and repeated handling on stress response and brain monoaminergic neurotransmitters in Senegalese sole (*Solea senegalensis*) juveniles. *Comp. Biochem. Physiol. – A Molecul. Integra. Physiol.*, 161: 18–26.
- Dabrowski K., Ciereszko A. (2001). Ascorbic acid and reproduction in fish: Endocrine regulation and gamete quality. *Aquacul. Res.*, 32: 623–638.
- Dediu L., Docan A., Crețu M., Grecu I., Mogodan A., Maoreanu M., Oprea L. (2021). Effects of Stocking Density on Growth Performance and Stress Responses of Bester and Bester ♀ × Beluga ♂ Juveniles in Recirculating Aquaculture Systems. *Animals*, 11: 2292.
- Do Huu H., Sang H.M., Thanh Thuy N.T. (2016). Dietary  $\beta$ -glucan improved growth performance, *Vibrio* counts, haematological parameters and stress resistance of pompano fish, *Trachinotus ovatus* Linnaeus, 1758. *Fish Shellfish Immunol.*, 54: 402–410.
- Dorts J., Silvestre F., Tu H.T., Tyberghein A.E., Phuong N.T., Kestemont P. (2009). Oxidative stress, protein carbonylation and heat shock proteins in the black tiger shrimp, *Penaeus monodon*, following exposure to endosulfan and deltamethrin. *Environ. Toxicol. Pharmacol.*, 28: 302–310.
- Ellis T., North B., Scott A.P., Bromage N.R., Porter M., Gadd D. (2002). The relationships between stocking density and welfare in farmed rainbow trout. *J. Fish Biol.*, 61: 493–531.
- Espelid S., Løkken G.B., Steiro K., Bøgwald J. (1996). Effects of cortisol and stress on the immune system in Atlantic Salmon (*Salmo salar* L.). *Fish Shellfish Immunol.*, 6: 95–110.
- Espina L., Somolinos M., Lorán S., Conchello P., García D., Pagán R. (2011). Chemical composition of commercial citrus fruit essential oils and evaluation of their antimicrobial activity acting alone or in combined processes. *Food Cont.*, 22: 896–902.
- Esteban M.Á., Rodríguez A., Ayala A.G., Meseguer J. (2004). Effects of high doses of cortisol on innate cellular immune response of seabream (*Sparus aurata* L.). *Gen. Comp. Endocrinol.*, 137: 89–98.
- Fajfer S., Meyers L., Willman G., Carpenter T., Hansen M.J. (1999). Growth of Juvenile Lake Sturgeon Reared in Tanks at Three Densities. *North Amer. J. Aquacul.*, 61: 331–335.
- Fatima S., Izhar S., Usman Z., Rashid F., Kanwal Z., Jabeen, Latif A. (2017). Effects of High Stocking Density on Condition Factor and Profile of Free Thyroxine and Cortisol in *Catla catla* (Hamilton, 1822) and *Labeo rohita* (Hamilton, 1822). *Turkish J. Fish. Aqua. Sci.*, 18.
- Gabr G.A., Ibrahim Y.S., Al-Shawi S.G., Abosaooda M., Gupta J., Oudaha K.H., Mavlonov K., Jalil A.T., Thalij K.M., Mustafa Y.F., Khodadadi M., Dadras M. (2023). Single or

- combined consumption of resveratrol and the probiotic, *Lactobacillus acidophilus* attenuate the effects of crowding stress on growth, immune characteristics, and antioxidant defense in the common carp, (*Cyprinus carpio*). *Aquacult. Rep.*, 29: 101471.
- Ganga R., Montero D., Bell J.G., Atalah E., Ganuza E., Vega-Orellana O., Tort L., Acerete L., Afonso J.M., Benitez-Sanatana T., Fernández Vaquero A., Izquierdo M. (2011). Stress response in sea bream (*Sparus aurata*) held under crowded conditions and fed diets containing linseed and/or soybean oil. *Aquaculture*, 311: 215–223.
- Gao Y., He Z., Vector H., Zhao B., Li Z., He J., Lee J.Y., Chu Z. (2017). Effect of stocking density on growth, oxidative stress and HSP 70 of pacific white Shrimp *Litopenaeus vannamei*. *Turkish J. Fish. Aqua. Sci.*, 17: 877–884.
- García Beltrán J.M., Espinosa C., Guardiola F.A., Esteban M.Á. (2017). Dietary dehydrated lemon peel improves the immune but not the antioxidant status of gilthead seabream (*Sparus aurata* L.). *Fish Shellfish Immunol.*, 64: 426–436.
- García de La Banda I., Lobo C., León-Rubio J.M., Tapia-Paniagua S., Balebona M.C., Moriñigo M.A., Moreno-Ventas X., Lucas L.M., Linares F., Arce F., Arijó S. (2010). Influence of two closely related probiotics on juvenile Senegalese sole (*Solea senegalensis*, Kaup 1858) performance and protection against *Photobacterium damsela* subsp. *piscicida*. *Aquaculture*, 306: 281–288.
- García F., Romera D.M., Gozi K.S., Onaka E.M., Fonseca F.S., Schalch S.H.C., Candeira P.G., Guerra L.O.M., Carmo F.J., Carneiro D.J., Martins M.I.E.G., Portella M.C. (2013). Stocking density of Nile tilapia in cages placed in a hydroelectric reservoir. *Aquaculture*, 410–411: 51–56.
- Gatesoupe F.J. (1999). The use of probiotics in aquaculture. *Aquaculture*, 180: 147–165.
- Ghafarifarsani H., Hoseinifar S.H., Adhami B., Rohani M.F., Van Doan H. (2023). Dietary gallic acid influences serum enzymatic parameters and immunological responses in *Cyprinus carpio* exposed to crowding stress. *Aquacult. Rep.*, 30: 101630.
- Gibtan A., Getahun A., Mengistou S. (2008). Effect of stocking density on the growth performance and yield of Nile tilapia [*Oreochromis niloticus* (L., 1758)] in a cage culture system in Lake Kuriftu, Ethiopia. *Aquacult. Res.*, 39: 1450–1460.
- Goh K.W., Abdul Kari Z., Wee, W., Van Doan H., Reduan M.F.H., Kabir M.A., Khoo M.I., Al-Amsyar S.M., Seong Wei L. (2023). The Roles of Polysaccharides in Carp Farming: A Review. *Animals*, 13: 244.
- Gómez G.D., Balcázar J.L. (2008). A review on the interactions between gut microbiota and innate immunity of fish. *FEMS Immunol. Medic. Microbiol.*, 52: 145–154.
- Güroy B., Şahin I., Mantoğlu S., Kayali S. (2012). Spirulina as a natural carotenoid source on growth, pigmentation and reproductive performance of yellow tail cichlid *Pseudotropheus acei*. *Aquacul. Int.*, 20: 869–878.
- Hamid N.K.A., Somdare P.O., Md Harashid K.A., Othman N.A., Kari Z.A., Wei L.S., Dawood M.A.O. (2022). Effect of papaya (*Carica papaya*) leaf extract as dietary growth promoter supplement in red hybrid tilapia (*Oreochromis mossambicus* × *Oreochromis niloticus*) diet. *Saudi J. Biol. Sci.*, 29: 3911–3917.
- Harikrishnan R., Thamizharasan S., Devi G., Van Doan H., Ajith Kumar T.T., Hoseinifar S.H., Balasundaram C. (2020). Dried lemon peel enriched diet improves antioxidant activity,

- immune response and modulates immuno-antioxidant genes in *Labeo rohita* against *Aeromonas sorbia*. *Fish Shellfish Immunol.*, 106: 675–684.
- Henrique M.M.F., Gomes E.F., Gouillou-Coustans M.F., Oliva-Teles A., Davies S.J. (1998). Influence of supplementation of practical diets with vitamin C on growth and response to hypoxic stress of seabream, *Sparus aurata*. *Aquaculture*, 161: 415–426.
- Herrera M., Mancera J.M., Costas B. (2019). The Use of Dietary Additives in Fish Stress Mitigation: Comparative Endocrine and Physiological Responses. *Front. Endocrinol.*, 10.
- Heshmati J., Farsi F., Shokri F., Rezaeinejad M., Almasi-Hashiani A., Vesali S., Sepidarkish M. (2018). A systematic review and meta-analysis of the probiotics and synbiotics effects on oxidative stress. *J. Funct. Food.*, 46: 66–84.
- Hoseini S.M., Abtahi B., Yousefi M. (2019). Effects of fasting on metabolic and immunological responses of common carp (*Cyprinus carpio*) to a further acute stress. *Aquacult. Res.*, 50: 1177–1185.
- Hoseini S.M., Yousefi M., Hoseinifar S.H., Van Doan H. (2019). Effects of dietary arginine supplementation on growth, biochemical, and immunological responses of common carp (*Cyprinus carpio* L.), stressed by stocking density. *Aquaculture*, 503: 452–459.
- Hoseinifar S.H., Hosseini M., Paknejad H., Safari R., Jafar A., Yousefi M., Van Doan H., Torfi Mozanzadeh M. (2019). Enhanced mucosal immune responses, immune related genes and growth performance in common carp (*Cyprinus carpio*) juveniles fed dietary *Pediococcus acidilactici* MA18/5M and raffinose. *Develop. Comp. Immunol.*, 94: 59–65.
- Hoseinifar S.H., Yousefi S., Van Doan H., Ashouri G., Gioacchini G., Maradonna F., Carnevali O. (2020). Oxidative Stress and Antioxidant Defense in Fish: The Implications of Probiotic, Prebiotic, and Synbiotics. *Rev. Fish. Sci. Aquacult.*, 29: 198–217.
- Hosseini S. A., Hoseini S.M. (2012). Effect of acute crowding stress on subsequent osmotic challenge and recovery in juvenile common carp *Cyprinus carpio* (Linnaeus). *Comp. Clin. Pathol.*, 21: 583–588.
- Hwang J.H., Lee S.W., Rha S.J., Yoon H.S., Park E.S., Han K.H., Kim S.J. (2013). Dietary green tea extract improves growth performance, body composition, and stress recovery in the juvenile black rockfish, *Sebastes schlegeli*. *Aquacult. Int.*, 21: 525–538.
- Iwama G.K., Vijayan M.M., Forsyth R.B., Ackerman P.A. (2015). Heat Shock Proteins and Physiological Stress in Fish. *Amer. Zool.*, 39: 901–909.
- Jeney G., Jeney Z. (2002). Application of immunostimulants for modulation of the non-specific defense mechanisms in sturgeon hybrid: *Acipenser ruthenus* × *A. baerii*. *J. Appl. Ichthyol.*, 18: 416–419.
- Jia R., Liu B.L., Feng W.R., Han C., Huang B., Lei J.L. (2016). Stress and immune responses in skin of turbot (*Scophthalmus maximus*) under different stocking densities. *Fish Shellfish Immunol.*, 55: 131–139.
- Jian J., Wu Z. (2004). Influences of traditional Chinese medicine on non-specific immunity of Jian Carp (*Cyprinus carpio* var. Jian). *Fish Shellfish Immunol.*, 16: 185–191.
- Junqueira V.B.C., Barros S.B.M., Chan S.S., Rodrigues L., Giavarotti L., Abud R.L., Deucher G.P. (2004). Aging and oxidative stress. *Mol. Asp. Med.*, 25: 5–16.

- Kari Z.A., Wee W., Abdul Hamid N.K., Dawood M.A.O., Azwanida Binti Zakaria N. N., & Wei, L. S. (2024). The roles of polysaccharides in tilapia farming: A review. *Aquacult. Fish.*, 9: 20–27.
- Kari Z. A., Wee W., Hamid N.K.A., Mat K., Rusli N.D., Khalid H.N.M., Sukri S.A.M., Harun H.C., Dawood M.A.O., Hakim A.H., Khoo M.I., Abd El-Razek I.M., Goh K.W., Wei L.S. (2022). Recent Advances of Phytobiotic Utilization in Carp Farming: A Review. *Aquacul. Nutr.*, 7626675.
- Karnatak G., Das B.K., Mishal P., Tayung T., Kumari S., Sarkar U.K., Das A.K., Ali Y. (2021). Impact of stocking density on growth, feed utilisation and survival of cage reared minor carp, *Labeo bata* (Hamilton, 1822) in Maithon reservoir, India. *Aquaculture*, 532: 736078.
- Kaur G., Athar M., Alam M.S. (2018). Pulegone: a potential candidate for the development of antimicrobial agents. *Biomed. Pharmacother.*, 107: 372–380.
- Kirihara N., Kamitomo M., Tabira T., Hashimoto T., Taniguchi H., Maeda T. (2018). Effect of probiotics on perinatal outcome in patients at high risk of preterm birth. *J. Obs. Gynaecol. Res.*, 44: 241–247.
- Kleinert M., Sachs S., Habegger K. M., Hofmann S.M. Müller T.D. (2019). Glucagon Regulation of Energy Expenditure. *Int. J. Mol. Sci.*, 20: 5407.
- Lambert Y., Dutil J.D. (2001). Food intake and growth of adult Atlantic cod (*Gadus morhua* L.) reared under different conditions of stocking density, feeding frequency and size-grading. *Aquacul.*, 192: 233–247.
- Lee J.S., Park S.A., Chung D., Lee H.G. (2011). Encapsulation of astaxanthin-rich Xanthophyllomyces dendrorhous for antioxidant delivery. *Int. J. Biol. Macromol.*, 49: 268–273.
- Lee Y.J., Corry P.M. (1998). Metabolic oxidative stress-induced HSP70 gene expression is mediated through SAPK pathway: Role of Bcl-2 and c-Jun NH2-terminal kinase. *J. Biol. Chem.*, 273: 29857–29863.
- Lewis S., Handy R.D., Cordi B., Billingham Z., Depledge M.H. (1999). Stress proteins (HSP's): Methods of detection and their use as an environmental biomarker. *Ecotoxicology*, 8: 351–368.
- Li W., Li D., Yang Q., Liu L., Liu J., Lu J., Wang Y., Tang R., Li L., Zhang X. (2023). Long-term crowding stress induces chronic inflammatory response and declines the immunity of grass carp (*Ctenopharyngodon idella*). *Aquaculture*, 577: 739976.
- Lin W., Li L., Chen J., Li D., Hou J., Guo H., Shen J. (2018). Long-term crowding stress causes compromised nonspecific immunity and increases apoptosis of spleen in grass carp (*Ctenopharyngodon idella*). *Fish Shellfish Immunol.*, 80: 540–545.
- Lindquist S., Craig E.A. (1988). The heat-shock proteins. *Ann. Rev. Gen.*, 22: 631–677.
- Liu B., Xu P., Xie J., Ge X., Xia S., Song C., Zhou Q., Miao L., Ren M., Pan L., Chen R. (2014). Effects of emodin and vitamin E on the growth and crowding stress of Wuchang bream (*Megalobrama amblycephala*). *Fish Shellfish Immunol.*, 40: 595–602.
- Liu H., Xu K., Xu M. (2004). Novel ECN-based congestion control algorithm with forecasting and verifying. *High Technol. Let.*, 14: 16.

- Liu J., Zhang W., Du X., Jiang J., Wang C., Wang X., Zhang Q., He Y. (2016). Molecular characterisation and functional analysis of the GATA4 in tongue sole (*Cynoglossus semilaevis*). *Comp. Biochem. Physiol. Part B: Biochem. Mol. Biol.*, 193: 1–8.
- Long L., Zhang H., Ni Q., Liu H., Wu F., Wang X. (2019). Effects of stocking density on growth, stress, and immune responses of juvenile Chinese sturgeon (*Acipenser sinensis*) in a recirculating aquaculture system. *Comp. Biochem. Physiol. Part C: Toxicol. Pharmacol.*, 219: 25–34.
- Loto I., Gutiérrez M.S., Barahona S., Sepúlveda D., Martínez-Moya P., Baeza M., Cifuentes V., Alcaíno J. (2012). Enhancement of carotenoid production by disrupting the C22-sterol desaturase gene (CYP61) in *Xanthophyllomyces dendrorhous*. *BMC Microbiol.*, 12: 235.
- Malviya, S., Jha, A., & Hettiarachchy, N. (2014). Antioxidant and antibacterial potential of pomegranate peel extracts. *J. Food Sci. Technol.*, 51: 4132–4137.
- Mani R., Natesan V. (2018). Chrysin: Sources, beneficial pharmacological activities, and molecular mechanism of action. *Phytochem.*, 145: 187–196.
- Martinez-Moya P., Niehaus K., Alcaíno J., Baeza M., Cifuentes V. (2015). Proteomic and metabolomic analysis of the carotenogenic yeast *Xanthophyllomyces dendrorhous* using different carbon sources. *BMC Genom.*, 16: 289.
- Mauri I., Romero A., Acerete L., MacKenzie S., Roher N., Callol A., Cano I., Alvarez M.C., Tort L. (2011). Changes in complement responses in Gilthead seabream (*Sparus aurata*) and European seabass (*Dicentrarchus labrax*) under crowding stress, plus viral and bacterial challenges. *Fish Shellfish Immunol.*, 30: 182–188.
- Mazeaud M.M., Mazeaud F., Donaldson E.M. (1977). Primary and Secondary Effects of Stress in Fish: Some New Data with a General Review. *Trans. Amer. Fish. Soc.*, 106: 201–212.
- Mohler J.W., King M.K., Farrell P.R. (2000). Growth and Survival of First-Feeding and Fingerling Atlantic Sturgeon under Culture Conditions. *North American Journal Aquacult.*, 62: 174–183.
- Mommsen T.P., Vijayan, M.M., Moon, T.W. (1999). Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. *Rev Fish Biol. Fish.*, 9: 211–268.
- Monir W., Abdel-Rahman M.A., El-Din Hassan S., Mansour E.S., Awad S.M.M. (2020). Pomegranate peel and moringa-based diets enhanced biochemical and immune parameters of Nile tilapia against bacterial infection by *Aeromonas hydrophila*. *Microb. Pathol.*, 145: 104202.
- Montero D., Marrero M., Izquierdo M.S., Robaina L., Vergara J.M., Tort L. (1999). Effect of vitamin E and C dietary supplementation on some immune parameters of gilthead seabream (*Sparus aurata*) juveniles subjected to crowding stress. *Aquaculture*, 171: 269–278.
- Morgan A.L., Thompson K.D., Auchinachie N.A., Migaud H. (2008). The effect of seasonality on normal haematological and innate immune parameters of rainbow trout *Oncorhynchus mykiss* L. *Fish Shellfish Immunol.*, 25: 791–799.
- Ngugi C.C., Oyoo-Okoth E., Muchiri M. (2017). Effects of dietary levels of essential oil (EO) extract from bitter lemon (*Citrus limon*) fruit peels on growth, biochemical, haemato-

- immunological parameters and disease resistance in Juvenile *Labeo victorianus* fingerlings challenged with *Aeromonas hydrophila*. *Aquacult. Res.*, 48: 2253–2265.
- Ngugi C.C., Oyoo-Okoth E., Mugo-Bundi J., Orina P.S., Chemoiwa E.J., Aloo P.A. (2015). Effects of dietary administration of stinging nettle (*Urtica dioica*) on the growth performance, biochemical, hematological and immunological parameters in juvenile and adult Victoria Labeo (*Labeo victorianus*) challenged with *Aeromonas hydrophila*. *Fish Shellfish Immunol.*, 44: 533–541.
- North B.P., Turnbull J.F., Ellis T., Porter M.J., Migaud H., Bron J., Bromage N.R. (2006). The impact of stocking density on the welfare of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 255: 466–479.
- Osuna-Jiménez I., Williams T.D., Prieto-Álamo M.J., Abril N., Chipman J.K., Pueyo C. (2009). Immune- and stress-related transcriptomic responses of *Solea senegalensis* stimulated with lipopolysaccharide and copper sulphate using heterologous cDNA microarrays. *Fish Shellfish Immunol.*, 26: 699–706.
- Pahor-Filho E., Castillo A.S.C., Pereira N.L., Pilarski F., Urbinati E.C. (2017). Levamisole enhances the innate immune response and prevents increased cortisol levels in stressed pacu (*Piaractus mesopotamicus*). *Fish Shellfish Immunol.*, 65: 96–102.
- Paray B.A., Hoseini S.M., Hoseinifar S.H., Van Doan H. (2020). Effects of dietary oak (*Quercus castaneifolia*) leaf extract on growth, antioxidant, and immune characteristics and responses to crowding stress in common carp (*Cyprinus carpio*). *Aquaculture*, 524: 735276.
- Pascoli F., Lanzano G.S., Negrato E., Poltronieri C., Trocino A., Radaelli G., Bertotto D. (2011). Seasonal effects on hematological and innate immune parameters in sea bass *Dicentrarchus labrax*. *Fish Shellfish Immunol.*, 31: 1081–1087.
- Peng C.H., Su J.D., Chyau C.C., Sung T.Y., Ho S.S., Peng C.C., Peng R.Y. (2007). Supercritical fluid extracts of rosemary leaves exhibit potent anti-inflammation and anti-tumor effects. *Biosci., Biotechnol. Biochem.*, 71: 2223–2232.
- Prieto-Álamo M. J., Abril N., Osuna-Jiménez I., Pueyo C. (2009). *Solea senegalensis* genes responding to lipopolysaccharide and copper sulphate challenges: Large-scale identification by suppression subtractive hybridisation and absolute quantification of transcriptional profiles by real-time RT-PCR. *Aqua. Toxicol.*, 91: 312–319.
- Rama S., Manjabhat S.N. (2014). Protective effect of shrimp carotenoids against ammonia stress in common carp, *Cyprinus carpio*. *Ecotoxicol. Environ. Safety.*, 107: 207–213.
- Reyes-Cerpa S., Vallejos-Vidal E., Gonzalez-Bown M.J., Morales-Reyes J., Pérez-Stuardo D., Vargas D., Imarai M., Cifuentes V., Spencer E., Sandino A.M., Reyes-López F.E. (2018). Effect of yeast (*Xanthophyllomyces dendrorhous*) and plant (Saint John's wort, lemon balm, and rosemary) extract based functional diets on antioxidant and immune status of Atlantic salmon (*Salmo salar*) subjected to crowding stress. *Fish. Shellfish. Immunol.*, 74: 250–259.
- Ringø E., Zhou Z., Vecino J.L.G., Wadsworth S., Romero J., Krogdahl Å., Olsen R.E., Dimitroglou A., Foey A., Davies S., Owen M., Lauzon H.L., Martinsen L.L., De Schryver P., Bossier P., Sperstad S., Merrifield D.L. (2016). Effect of dietary components on the gut microbiota of aquatic animals. A never-ending story?. *Aquacult. Nutr.*, 22: 219–282.

- Rotruck J. ., Pope A.L., Ganther H.E., Swanson A.B., Hafeman D.G., Hoekstra W.G. (1973). Selenium: biochemical role as a component of glutathione peroxidase. *Science*, 179: 588–590.
- Sadeghi F., Ahmadifar E., Shahriari Moghadam M., Ghiyasi M., Dawood M.A.O., Yilmaz S. (2021). Lemon, *Citrus aurantifolia*, peel and *Bacillus licheniformis* protected common carp, *Cyprinus carpio*, from *Aeromonas hydrophila* infection by improving the humoral and skin mucosal immunity, and antioxidative responses. *J. World Aquacult. Soc.*, 52: 124–137.
- Sadoul B., Vijayan M.M. (2016). Stress and Growth. In: *Fish Physiology*. Vol. 35, pp. 167–205.
- Saillant E., Fostier A., Haffray P., Menu B., Laureau S., Thimonier J., Chatain B. (2003). Effects of rearing density, size grading and parental factors on sex ratios of the sea bass (*Dicentrarchus labrax* L.) in intensive aquaculture. *Aquaculture*, 221: 183–206.
- Salas-Leiton E., Anguis V., Manchad, M., Cañavate J. P. (2008). Growth, feeding and oxygen consumption of Senegalese sole (*Solea senegalensis*) juveniles stocked at different densities. *Aquaculture*, 285: 84–89.
- Salomão R.A.S., De Paula T.G., Zanella B.T.T., Carvalho P.L.P.F., da Silva Duran B.O., Valente J.S., de Almeida Fantinatti B.E., Fernandes A.A., Barros M.M., Mareco E.A., Carvalho R.F., dos Santos V.B., Dal-Pai-Silva M. (2019). The combination of resveratrol and exercise enhances muscle growth characteristics in pacu (*Piaractus mesopotamicus*). *Comp. Biochem. Physiol. Part A : Molecul. Integrat. Physiol.*, 235: 46–55.
- Sanders B.M. (1993). Stress proteins in aquatic organisms: An environmental perspective. *Critical Rev. Toxicol.*, 23: 49–75.
- Schreck C. B. (2010). Stress and fish reproduction: The roles of allostasis and hormesis. *Gen. Comp. Endocrinol.*, 165: 549–556.
- Schreck C.B., Tort L. (2016). The Concept of Stress in Fish. In: *Fish Physiology*. Vol. 35, pp. 1–34.
- Seong Wei L., Mohamad Sukri S.A., Tahiluddin A.B., Abdul Kari Z., Wee W., Kabir M.A. (2024). Exploring beneficial effects of phytobiotics in marine shrimp farming: A review. *Heliyon*, 10: e31074.
- Sharma V., Sharma A., Kansal L. (2010). The effect of oral administration of *Allium sativum* extracts on lead nitrate induced toxicity in male mice. *Food Chem. Toxicol.*, 48: 928–936.
- Singh B., Singh J.P., Kaur A., Singh N. (2019). Antimicrobial potential of pomegranate peel: a review. *Int. J. Food Sci. Technol.*, 54: 959–965.
- Stompor-gorący M., Bajek-bil A., Machaczka M. (2021). Chrysin: Perspectives on contemporary status and future possibilities as pro-health agent. *Nutrients.*, 13: 2038.
- Sukri S.A.M., Andu Y., Sarijan S., Khalid H.N.M., Kari Z.A., Harun H.C., Rusli N.D., Mat K., Khalif R.I.A.R., Wei L.S., Rahman M.M., Hakim A.H., Lokman N.H.N., Hamid N.K.A., Khoo M.I., Doan H.V. (2023). Pineapple waste in animal feed: A review of nutritional potential, impact and prospects. *Ann. Anim. Sci.*, 23: 339–352.

- Takanashi S., Miura A., Abe K., Uchida J., Itoi S., Sugita H. (2014). Variations in bile tolerance among *Lactococcus lactis* strains derived from different sources. *Folia Microbiol.*, 59: 289–293.
- Tan C., Zhou H., Wang X., Mai K., He G. (2019). Resveratrol attenuates oxidative stress and inflammatory response in turbot fed with soybean meal based diet. *Fish. Shellfish. Immunol.*, 91: 130–135.
- Tapia-Paniagua S.T., Chabrilón M., Díaz-Rosales P., de la Banda I.G., Lobo C., Balebona M.C., Moriñigo M.A. (2010). Intestinal Microbiota Diversity of the Flat Fish *Solea senegalensis* (Kaup, 1858) Following Probiotic Administration. *Microb. Ecol.*, 60: 310–319.
- Tapia-Paniagua S.T., Vidal S., Lobo C., Prieto-Álamo M.J., Jurado J., Cordero H., Cerezuela R., García de la Banda I., Esteban M.A., Balebona M.C., Moriñigo M.A. (2014). The treatment with the probiotic *Shewanella putrefaciens* Pdp11 of specimens of *Solea senegalensis* exposed to high stocking densities to enhance their resistance to disease. *Fish Shellfish Immunol.*, 41: 209–221.
- Tort L. (2011). Stress and immune modulation in fish. *Develop. Comp. Immunol.*, 35: 1366–1375.
- Tort L., Rotllant J., Rovira L. (1998). Immunological suppression in gilthead sea bream *Sparus aurata* of the North-West Mediterranean at low temperatures. *Comp. Biochem. Physiol. Part A: Molecul. Integra. Physiol.*, 120: 175–179.
- Tungmunnithum D., Thongboonyou A., Pholboon A., Yangsabai A. (2018). Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview. *Medicines*, 5.
- Wang H., Lu T., Jiang P. (2014). Mathematical model and numerical simulation of biological porous medium during hot air drying. *Transact. Chin. Soc. Agricult. Engin.*, 30: 325–333.
- Wee W., Abdul Hamid N.K., Mat K., Khalif R.I.A.R., Rusli N.D., Rahman M.M., Kabir M.A., Wei L.S. (2024). The effects of mixed prebiotics in aquaculture: A review. *Aquacult. Fish.*, 9: 28–34.
- Wei, L. S., Goh, K. W., Abdul Hamid, N. K., Abdul Kari, Z., Wee, W., & Van Doan, H. (2022). A mini-review on co-supplementation of probiotics and medicinal herbs: Application in aquaculture. *Front. Vet. Sci.*, 9.
- Wilson W.N., Baumgarner B.L., Watanabe W.O., Alam M.S., Kinsey S.T. (2015). Effects of resveratrol on growth and skeletal muscle physiology of juvenile southern flounder. *Comp. Biochem. Physiol. -Part A : Molecul. Integ. Physiol.*, 183: 27–35.
- Xi W., Lu J., Qun J., Jiao B. (2017). Characterisation of phenolic profile and antioxidant capacity of different fruit part from lemon (*Citrus limon* Burm.) cultivars. *J. Food Sci. Technol.*, 54: 1108–1118.
- Xie J., Liu B., Zhou Q., Su Y., He Y., Pan L., Ge X., Xu P. (2008). Effects of anthraquinone extract from rhubarb *Rheum officinale* Bail on the crowding stress response and growth of common carp *Cyprinus carpio* var. Jian. *Aquaculture*, 281: 5–11.
- Yengkokpam S., Debnath D., Sahu N.P., Pal A.K., Jain K.K., Baruah K. (2016). Dietary protein enhances non-specific immunity, antioxidative capability and resistance to *Aeromonas*

- hydrophila* in *Labeo rohita* fingerlings pre-exposed to short feed deprivation stress. Fish. Shellfish. Immunol., 59: 439–446.
- Yin G., Jeney G., Racz T., Xu P., Jun X., Jeney Z. (2006). Effect of two Chinese herbs (*Astragalus radix* and) on non-specific immune response of tilapia, *Oreochromis niloticus*. Aquaculture, 253: 39–47.
- Yousefi M., Adineh H., Ghadamkheir M., Hashemianfar S.A.M., Yilmaz S. (2023). Effects of dietary Pennyroyal essential oil on growth performance, digestive enzymes' activity, and stress responses of common carp, *Cyprinus carpio*. Aquacult. Rep., 30: 101574.
- Yousefi M., Hoseini S.M., Kulikov E.V., Babichev N.V., Bolshakova M.V., Shopinskaya M.I., Rogov R.V., Zharov A.N. (2023). Effects of dietary pomegranate peel supplementation on growth performance and biochemical responses of common carp, *Cyprinus carpio*, to chronic crowding stress. Aquacult. Rep., 30: 101532.
- Yousefi M., Hoseini S.M., Vatnikov Y.A., Kulikov E.V., Drukovsky S.G. (2019). Rosemary leaf powder improved growth performance, immune and antioxidant parameters, and crowding stress responses in common carp (*Cyprinus carpio*) fingerlings. Aquaculture, 505: 473–480.
- Yousefi M., Nedaei S., Naderi Farsani M., Ghafarifarsani H., Zhang M.L., Du Z.Y. (2023). Dietary chrysin supplementation improves growth performance, immune responses, antioxidant status, and resistance against crowding stress in rainbow trout. Aquacult. Rep., 32: 101708.
- Yousefi M., Shabunin S.V., Vatnikov Y.A., Kulikov E.V., Adineh H., Khademi Hamidi M., Hoseini S.M. (2020). Effects of lavender (*Lavandula angustifolia*) extract inclusion in diet on growth performance, innate immunity, immune-related gene expression, and stress response of common carp, *Cyprinus carpio*. Aquaculture, 515: 734588.
- Zeinali M., Rezaee S. A., Hosseinzadeh H. (2017). An overview on immunoregulatory and anti-inflammatory properties of chrysin and flavonoids substances. Biomed. Pharmacother., 92: 998–1009.
- Zhang C.N., Li X.F., Xu W.N., Jiang G.Z., Lu K.L., Wang L.N., Liu W.B. (2013). Combined effects of dietary fructooligosaccharide and *Bacillus licheniformis* on innate immunity, antioxidant capability and disease resistance of triangular bream (*Megalobrama terminalis*). Fish. Shellfish. Immunol., 35: 1380–1386.
- Zheleva-Dimitrova D., Nedialkov P., Kitanov G. (2010). Radical scavenging and antioxidant activities of methanolic extracts from *Hypericum* species growing in Bulgaria. Pharmacog. Mag., 6: 74–78.

Received: 28 X 2024

Accepted: 13 III 2025